



FLURON[®] GOLD POUR ON

Product information

CONTENTS

1 	Name of the veterinary medicinal product	3
2 	Qualitative and quantitative composition	3
3 	Pharmaceutical form	3
4 	Indications	3
5 	Administration and posology	3
6 	Interval between applications	4
7 	Contraindications and limitations of use	4
8 	Adverse reactions	5
9 	Precautions	5
10 	Withdrawal periods	6
11 	Storage conditions	6
12 	Presentations	6
13 	Shelf life	6

These pages are general information pages. No guarantee is given as to the completeness of the information contained or its compliance with national regulatory requirements. Users should consult the local site of their countries to obtain information that complies with applicable national regulations.

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Fluron Gold Pour On.

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each 100 ml contains:

Cypermethrin	5.0 g
Chlorpyrifos	7.0 g
Fluazuron	2.5 g
Piperonyl butoxide	5.0 g
Excipients q.s.	100,00mL

3. PHARMACEUTICAL FORM

Pour-on solution.

4. INDICATIONS

Fluron Gold Pour On is an ectoparasiticide for topical application (pour on), containing cypermethrin 5%, chlorpyrifos 7%, fluazuron 2.5% and piperonyl butoxide, indicated for the treatment of the main bovine ectoparasites, namely:

- Treatment of infestations caused by eggs, larvae, nymphs and adults stages of the tick *Rhipicephalus microplus*;
- Treatment of infestations caused by larvae of *Dermatobia hominis* (warble);
- Treatment of infestations caused by *Haematobia irritans* (hornfly);

The product can be used in pregnant females at any stage of pregnancy.

5. ADMINISTRATION AND POSOLOGY

Administration and dosage

Posology:

Fluron Gold Pour On should be applied at a dose of 1mL for every 10kg of live bodyweight.

The table below shows the dosing scheme of the product:

Weight of the animal	Fluron Gold Pour On dosage
100 kg	10 mL
150 kg	15 mL
200 kg	20 mL
250 kg	25 mL
300 kg	30 mL
350 kg	35 mL
400 kg	40 mL
450 kg	45 mL
500 kg	50 mL

Note: In animals over 500 kg, use the dosage of 1mL for every 10kg of live bodyweight.

How to use:

Fluron Gold Pour On should be applied topically, along the backline of the animal homogeneously, from the neck to the tail.

6. INTERVAL BETWEEN APLICATIONS

Studies with the veterinary medicinal product show that anti-tick efficacy is maintained for up to 110 days, efficacy against warble is maintained for 7 days and the effectiveness against hornfly is maintained for up to 24 days after treatment.

7. CONTRAINDICATIONS AND LIMITATIONS OF USE

Contraindications:

Do not use the veterinary medicinal product on animals sensitive to the active ingredients.

Do not use the veterinary medicinal product on animals at the hottest hours of the day.

Do not use the veterinary medicinal product on rainy days.

Do not use the veterinary medicinal product in other indications apart from the ones mentioned in the leaflet.

Limitations of Use:

Do not use the veterinary medicinal product in weak, sick or animals or with hepatic conditions.

Do not use the product in cattle producing milk for human consumption.

The veterinary medicinal product safety has not been evaluated in animals under 18 months.

Some animals may have sensitivity to the product such as: head and tail swing, agitation and decreased appetite, but these symptoms tend to disappear 24 hours after the application of the product.

In more severe cases of hypersensitivity due to incorrect use of the product, such as the non-homogeneous distribution of the product along the back of the animal or overdose, the animal may have skin reactions.

8. ADVERSE REACTIONS

If you detect any serious side effect or other effects not mentioned in the leaflet, please report immediately to the veterinarian.

9. PRECAUTIONS

- Respect the dosages and recommendations of use of the veterinary medicinal product.
- Do not ingest.
- Wear gloves during handling to avoid contact with the skin.
- Do not smoke or eat during the handling of the veterinary medicinal product.
- Always wash hands after handling any veterinary product.
- In case of accidental contact with skin, eyes and mouth, rinse thoroughly with water.
- Do not store close to food, beverages, other medicines, toiletries or household care products.
- Keep this or any other veterinary medicinal product out of the reach of children and domestic animals.
- Do not re-use empty packaging and dispose them in accordance with current legislation to avoid contamination of the environment.
- Use eye protector. The veterinary medicinal product is irritating to the eyes. If the veterinary medicinal product contacts the eyes, wash them immediately.
- Wear rubber gloves, overalls, long sleeves, waterproof apron and boots. Irritating and/or absorbable product by the skin. If the veterinary medicinal product contacts with the skin, wash it immediately.

First aid:

In case of contact with skin and eyes: remove contaminated garments and wash it with plenty of running water for 15 minutes. In case of persistent irritation consult a doctor taking him the package leaflet of the veterinary medicinal product.

If ingested: do not induce vomiting and do not administer cathartics if the victim has diarrhea. Consult a doctor taking the package leaflet of the veterinary medicinal product.

Instructions for the doctor: support medication according to symptoms and their intensity.

Specific antidote: Atropine sulfate - 1 to 6 mg every 30 minutes, until mild atropinization (never administer atropine sulfate before the appearance of poisoning symptoms).

10. WITHDRAWAL PERIODS

Cattle:

Meat and offal: 56 days

Milk: this product must not be applied to cattle producing milk for human consumption.

The use of the veterinary medicinal product in conditions other than those indicated in this package leaflet may cause the presence of residues of the veterinary medicinal product above the approved limits, making the food of animal origin unfit for consumption.

11. STORAGE CONDITIONS

Store in a dry place, away from sunlight and at a temperature between 15 and 30 °C.

12. PRESENTATIONS

1L and 2L vials.

13. SHELF LIFE

Shelf-life of the veterinary medicinal product as packaged for sale: 2 years.

Shelf-life after first opening the immediate packaging: 6 months.

Sale under prescription and administration under the guidance of the veterinarian.