



VERIBEN[®]

Product information

CONTENTS

1 	Name of the veterinary medicinal product	3
2 	Composition	3
3 	Properties	3
4 	Indications	3
5 	Administration and dosage	3
6 	Contraindications	3
7 	Side effects	4
8 	Precautions	4
9 	Withdrawal Period	4
10 	Storage Conditions	4
11 	Presentations	4

These pages are general information pages. No guarantee is given as to the completeness of the information contained or its compliance with national regulatory requirements. Users should consult the local site of their countries to obtain information that complies with applicable national regulations.

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Veriben. Soluble granules for solution for injection. Protozoocide.

2. COMPOSITION

For the 2.36 g sachet:

Diminazene diacetate	1.05 g
Excipient (Antipyrine) q.s.	1 sachet (2.36 g)

For the 23.6 g sachet:

Diminazene diacetate	10.5 g
Excipient (Antipyrine) q.s.f	1 sachet (23.6 g)

3. PROPERTIES

The active ingredient of VERIBEN® is diminazene diacetate, drug of the diamidines family. Diminazene is highly active against both *Trypanosoma* and *Babesia* spp. It is also of value in the treatment of theileriosis due to *Theileria annulata*.

Diminazene has become the most commonly used therapeutic agent for trypanosomosis in animals.

The antipyrine, included in the excipient, ensures a complete solubility of the active ingredient and acts as an help to struggle against inflammation and fever.

4. INDICATIONS

Cattle, sheep, goats: Treatment of protozoan diseases:

- Trypanosomosis due to *Trypanosoma congolense*, *T. vivax* and *T. brucei*.
- Piroplasmosis due to *Babesia bovis*, *B. bigemina*, *B. ovis*, *B. motasi*.
- Theileriosis due to *Theileria annulata*.

5. ADMINISTRATION AND DOSAGE

Deep intramuscular route.

The recommended dosage is 3.5 mg of diminazene diacetate per kg bodyweight in a single injection. Do not exceed a total dose of 4 g of active ingredient in any animal.

Cattle, sheep, goats: Prepare the injectable solution by adding 15 ml of sterile water to each 2.36 g sachet. For the 23.6 g sachet it is necessary to add 150 ml of sterile water. Mix well water and powder before sucking the sterile solution into the syringe. Inject the prepared solution according to the following table:

Body weight	Volume of the reconstituted solution	As quantity of diminazene diaceturate
100 kg	5.8 ml	350 mg
200 kg	11.5 ml	700 mg
300 kg	17.3 ml	1050 mg
400 kg	23.0 ml	1400 mg

One injection is enough for the treatment.

6. CONTRAINDICATIONS

Do not administer to other species than those registered.

7. SIDE EFFECTS

Spermatic decrease may be produced sometimes on caprine after a daily administration of VERIBEN®.

8. PRECAUTIONS

In case of large volume to be injected, it is recommended to divide it into two separate sites. Keep out of reach of children.

9. WITHDRAWAL PERIOD

Meat: 21 days.

Milk: 3 days.

10. STORAGE CONDITIONS

Sealed sachets can be stored in a cool dry place below 30°C.

The reconstituted solution can be stored during 15 days in a fridge, protected from light and in a closed glass bottle.

11. PRESENTATIONS

Box with 10 x 2.36 g sachets

Box with 10 x 23.6 g sachets.

FOR VETERINARY USE ONLY

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.