



VERIDIUM[®]

Product information

CONTENTS

1 	Name of the veterinary medicinal product	3
2 	Composition	3
3 	Properties	3
4 	Indications	3
5 	Contraindications	3
6 	Side effects	3
7 	Administration and dosage	3
8 	Withdrawal Period	4
9 	Presentations	4
10 	Storage Conditions	4

These pages are general information pages. No guarantee is given as to the completeness of the information contained or its compliance with national regulatory requirements. Users should consult the local site of their countries to obtain information that complies with applicable national regulations.

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Veridium. Powder for injectable solution.

2. COMPOSITION

Sachet of 125 mg:

Isometamidium chloride hydrochloride 125 mg

Sachet of 1 g:

Isometamidium chloride hydrochloride 1 g

3. PROPERTIES

Isometamidium, an aminophenanthridium compound, has trypanocidal action used in curative and preventive treatment of animal trypanosomosis.

Its spectrum of activity ranges from *T. vivax*, *T. congolense* to *T. brucei* and *T. evansi*. It could be administered to the following target species: ruminants (cattle, goat, sheep) and camels.

4. INDICATIONS

Cattle (incl. zebu), sheep, goats, camels:

Curative and preventive treatment of trypanosomosis.

5. CONTRAINDICATIONS

Do not administer to other species as registered.

6. SIDE EFFECTS

Transitory discomfort may be observed in some animals following intramuscular injection. This reaction will disappear spontaneously without any treatment.

7. ADMINISTRATION AND DOSAGE

Cattle, sheep, goats:

Deep intramuscular route in the neck muscles.

- Sterile or distilled water should be used to prepare the solution which must be used extemporaneously or stored in the fridge for 24 hours maximum.

Camels: treatment of camels trypanosomosis: 0.5 mg/kg through slow intravenous injection following usual aseptic recommendations.

Do not inject VERIDIUM with another trypanocide at the same time.

Body weight	CURATIVE TREATMENT	PREVENTIVE TREATMENT
	Prepare a 2% solution by dissolving 1g in 50 mL of water	
100 kg	2.5 mL	5 mL
200 kg	5 mL	10 mL
300 kg	7.5 mL	15 mL
400 kg	10 mL	20 mL (2 sites)

8. WITHDRAWAL PERIOD

Meat: 1 month.

Milk: nil.

9. PRESENTATIONS

Box with 10 x 125 mg sachet.

Box with 10 x 1 g sachet.

10. STORAGE CONDITIONS

Store below 30°C in a dry place.