

FOR ANIMAL USE ONLY

2 4

Maxicare®

Reg. No. G3145 Act 36/1947,

Namibia Reg. No. V02/18.1.8/762 Act 13/2003



A roundworm, lungworm, milk tapeworm, liver fluke and nasal bot remedy with residual protection against re-infestation of wireworm and hookworm in sheep and goats.

CAUTION

**COMPOSITION: Albendazole 1,9 % m/v,
Closantel 3,0 % m/v**

Store in a cool place and protect from direct sunlight.

WARNINGS

- **NOT FOR USE IN MILK GOATS WHOSE MILK IS INTENDED FOR HUMAN CONSUMPTION.**
- Allow 30 days between last dosing and slaughter for human consumption.
- Management practices should always ensure that all drenching procedures are carefully carried out.
- It is particularly important not to exceed the recommended dose in ewes during the mating season, or during the first month of pregnancy.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.
- Keep out of reach of children, uninformed persons and animals.

PRECAUTIONS

- It is recommended that all sheep be vaccinated against Pulpy Kidney disease before dosing with any worm remedy.
- Do not treat animals more frequently than every three weeks.
- Wash hands thoroughly after use.
- Handle with care.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED.

Use as an oral drench. Shake well before use.

Dose rate for sheep and goats: 2,0 ml/10 kg body mass.

EFFICACY

Worm Species	Adult	Immature
Wireworm	*	*
Brown stomachworm	*	*
Bankruptworm	*	*
Long-necked bankruptworm	*	*
Hookworm	*	*
Nodular worm	*	*
Large-mouthed bowelworm	*	*
Lungworm	*	*
Liver fluke	*	** (6 weeks old)
Giant liver fluke	*	** (8 weeks old)
Milk tapeworm	Class1	

Definitions: * Controls = 90 % - 100 % effective.
** Aids in the control = 60 % - 89 % effective.

Class 1: 100 % effective in more than 80 % of treated animals.

Controls all stages of nasal bot.

Prevents roundworm eggs present in the animal at the time of dosing, from hatching.

DOSAGE TABLE

Body Mass (kg)	Dose (ml)	No. of doses per 200 ml
10	2,0	100
20	4,0	50
30	6,0	33
40	8,0	25
50	10,0	20

RESIDUAL EFFECT

Worm species	Interval between treatment and re-infestation
Wireworm	3 weeks*
Hookworm	3 weeks*

Definition:

* Controls = 90 % - 100 % effective.

② Benzimidazoles ④ Salicylanilides

Maxicare®

Reg. Nr. G3145 Wet 36/1947,
Namibiese Reg. Nr. V02/18.1.8/762. Wet 13/2003



'n Middel teen rondewurm, longwurm, melklintwurm, lewerslak en neusvliegjarwes (neuswurm) met nawerking teen herbesmetting van haarwurm en haakwurm in skape en bokke.

VERSIGTIG

SAMESTELLING: Albendasool 1,9 % m/v,
Klosantel 3,0 % m/v.

Berg in 'n koel plek en beskerm teen direkte sonlig.

WAARSKUWINGS

- **MOET NIE IN BOKKE GEBRUIK WAARVAN DIE MELK VIR MENSLIKE VERBRUIK BESTEM IS NIE.**
- Moet nie diere vir menslike verbruik binne 30 dae na behandeling slag nie.
- Volg bestuurspraktyke wat sal verseker dat dosering versigtig uitgevoer word.
- Dit is veral belangrik om nie die dosis te oorskry in ooe gedurende paartyd en die eerste maand van dragtigheid nie.
- Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.
- Hou buite bereik van kinders, ongeligte persone en diere.

VOORSORGMAATREËLS

- Dit word aanbeveel dat alle skape teen Bloednier siekte ingeënt word voordat dosering met enige wurmmiddel plaasvind.
- Moet nie diere meer dikwels as elke drie weke behandel nie.
- Was hande deeglik na gebruik.
- Hanteer versigtig.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANBEVEEL

Gebruik as 'n mondelingse dosering. Skud deeglik voor gebruik.

Dosis vir skape en bokke: 2,0 ml/10 kg liggaamsmassa.

DOELTREFFENDHEID

Wurmsort	Volwasse	Onvolwasse
Haarwurm	*	*
Bruinmaagwurm	*	*
Bankrotwurm	*	*
Langnekbankrotwurm	*	*
Haakwurm	*	*
Knoppieswurm	*	*
Grootbekwurm	*	*
Longwurm	*	*
Lewerslak	*	** (6 weke oud)
Reuse lewerslak	*	** (8 weke oud)
Melklintwurm	Klas 1	

Definisies: * Beheer = 90 % - 100 % doeltreffend.
** 'n Hulpmiddel in die beheer = 60 % - 89 % doeltreffend.

Klas 1: 100 % doeltreffend in meer as 80 % van behandelde diere.

Beheer alle stadia van neusvliegjarwes.

Verhoed dat rondewurmeiers, wat tydens dosering in die dier teenwoordig is, uitbroei.

DOSISTABEL

Liggaams-massa (kg)	Dosis (ml)	Aantal dosisse per 200 ml
10	2,0	100
20	4,0	50
30	6,0	33
40	8,0	25
50	10,0	20

NAWERKING

Wurmsort	Tydperk tussen behandeling en herbesmetting
Haarwurm	3 weke*
Haakwurm	3 weke*

Definisie:

*Beheer = 90 % - 100 % doeltreffend.