

APPROVED

By Nicky Scholtz at 9:46 am, Jun 18, 2021

APPROVED

By Thandiwe Mtinkulu at 12:10 pm, Jul 06, 2021

FOR ANIMAL USE ONLY

Tenaline® L.A.Reg. No. G2436 (Act 36/1947)
Namibia: [NSO] Reg. No. V97/17.1.2/1097 (Act 13/2003)**LONG ACTING**

For the treatment of tick-borne gallsickness (Anaplasmosis), heartwater, pneumonia, footrot, joint-ill, navel-ill and pink eye (Infectious Bovine Keratoconjunctivitis) in cattle, sheep and goats and for the treatment of pneumonia, joint-ill and navel-ill in pigs.

CAUTION**STORAGE INSTRUCTIONS**

Store in a cool dark place. Some darkening of the solution may occur without affecting the potency.

The product can be used for one month after breaching the container.

COMPOSITION

Oxytetracycline dihydrate 200 mg/mL.

WARNINGS

Animals may not be slaughtered for human consumption within 21 days of the last treatment.

Milk from treated cows may not be used for human consumption within 8 days of the last treatment.

Use only in cattle, sheep, goats and pigs.

Do not inject intravenously.

Do not inject piglets with iron preparations on the same day.

Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

Keep out of reach of children, uninformed persons and animals.

PRECAUTIONS

Consult your veterinarian for an accurate diagnosis.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

For the treatment of conditions caused by, or associated with oxytetracycline susceptible organisms, where prolonged antibiotic treatment is indicated, or where frequent animal handling and repeat treatment is not feasible.

Cattle: For the treatment of tick-borne gallsickness (anaplasmosis), heartwater, pneumonia, footrot, joint-ill, navel-ill and pink eye (Infectious Bovine Keratoconjunctivitis).

Sheep and goats: For the treatment of heartwater, pneumonia, footrot, joint-ill and navel-ill.

Pigs: For the treatment of pneumonia, joint-ill and navel-ill.

Do not inject intravenously.

The basic dosage rate is 1 mL per 10 kg body mass by deep intramuscular injection (except in piglets - see below).

The recommended site of injection is the side of the neck.

Usually a single treatment is sufficient, but a second injection may be given after 72 hours in more chronic cases (eg. Pink eye in cattle).

A temporary swelling may be seen at the site of injection.

DOSAGE

Cattle: 1 mL per 10 kg body mass by deep intramuscular injection. Maximum 10 mL per injection site.

Sheep and goats: 1 mL per 10 kg body mass by deep intramuscular injection. Maximum 5 mL per injection site.

Piglets under 2 kg body mass: 0.5 mL subcutaneously behind the ear.

Piglets 2 - 10 kg body mass: 1 mL subcutaneously behind the ear.

Larger pigs: 1 mL per 10 kg body mass by deep intramuscular injection. Maximum 10 mL per injection site.

PRESENTATION

Multi-dose vials containing 100 mL and 500 mL solution.

REGISTRATION HOLDER:

Ceva Animal Health (Pty) Ltd.

(Reg. No. 1973/016009/07) - P O Box 7707 - Halfway House - 1685

Tel. 011 312 4088

55209-05



LANGWE

APPROVED

By Nicky Scholtz at 9:46 am, Jun 18, 2021

Vir die behandeling van longontsteking, wrotpootjie, septiese gewrigsontsteking en naelstringontsteking by beeste, skape en bokke en vir die behandeling van longontsteking, septiese gewrigsontsteking en naelstringontsteking by varke.

VERSIGTIG

BERGINGSAAWYSINGS

Berg in 'n koel, donker plek. Die preparaat mag gedurende gebruik verdonker, maar die sterkte bly onveranderd.

Die produk kan gebruik word vir 1 maand na oopmaak.

SAMESTELLING

Oksitetrasiklien dihidraat 200 mg/ml.

WAARSKUWINGS

Moenie die behandelde diere binne 21 dae na die laaste behandeling vir menslike verbruik slag nie.

Melk van behandelde koeie moet nie vir menslike verbruik binne 8 dae na die laaste behandeling gebruik word nie.

Gebruik slegs by beeste, skape, bokke en varke.

Moet nie binnears toedien nie.

Moenie varkies op dieselfde dag met ysterpreparate inspuit nie.

Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

Hou buite bereik van kinders, oningeligte persone en diere.

VOORSORGMATREËLS

Raadpleeg u veearts vir 'n akkurate diagnose.

GEBRUIKSAAWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

Vir die behandeling van toestande veroorsaak deur, of geassosieer met oksitetrasikliengevoelige organismes, waar langdurige antibiotiese werking verlang word, of waar gedurige hantering en herhaalde behandeling van diere nie prakties is nie.

Beeste: vir die behandeling van bosluisoorgedraagde galsiekte (anaplasmosis), hartwater, longontsteking, wrotpootjie, septiese gewrigsontsteking, naelstringontsteking en pienkoog (Aansteeklike Beeskeratokonjunktivitis).

Skape en bokke: vir die behandeling van hartwater, longontsteking, wrotpootjie, gewrigsontsteking en naelstringontsteking.

Varke: vir die behandeling van longontsteking, septiese gewrigsontsteking, en naelstringontsteking.

Moet nie binnears inspuit nie.

Die basiese dosis is 1 ml per 10 kg liggaamsmassa deur diep binnespiers inspuiting (behalwe by jong varkies - sien onder).

Die aanbevole plek vir inspuiting is aan die kant van die nek.

'n Enkele inspuiting sal gewoonlik voldoende wees, maar behandeling kan na 72 uur herhaal word in meer kroniese gevalle (bv. pienkoog by beeste).

Tydlike gelokaliseerde swelsel by die plek van inspuiting mag waargeneem word.

DOSIS

Beeste: 1 ml per 10 kg liggaamsmassa deur diep binnespiers inspuiting. Maksimum van 10 ml per inspuitingsplek.

Skape en bokke: 1 ml per 10 kg liggaamsmassa deur diep binnespiers inspuiting. Maksimum van 5 ml per inspuitingsplek.

Varkies onder 2 kg liggaamsmassa: 0,5 ml onderhuids agter die oor toegedien.

Varkies 2 - 10 kg liggaamsmassa: 1 ml onderhuids agter die oor toegedien.

Groter varke: 1 ml per 10 kg liggaamsmassa deur diep binnespiers inspuiting. Maksimum van 10 ml per inspuitingsplek.

AANBIEDING

Meervoudige dosishouers wat 100 ml en 500 ml oplossing bevat.

REGISTRASIEHOUER:

Ceva Animal Health (Edms) Bpk.

(Reg. Nr. 1973/016009/07) - Posbus 7707 - Halfway House - 1685

Tel. 011 312 4088

NOT TENALINE LA. CLAS ZA
200X160/FP 35,5X160
RECTO/VERSO
CODE ARTICLE : 55209-05
BLACK
WARM RED

TENALINE LA CLAS AFRIQUE DU SUD
ID 70874
CORPS : 9 pts

LOUIS 07/06/21 - 17/06/21