

FOR ANIMAL USE ONLY

Consult a veterinarian before use

Ovipast

FOR SHEEP & GOATS

Reg. No. G4356 (Act 36 of 1947)

INDICATIONS

Ovipast is an adjuvanted multi-valent vaccine, without antibiotics or preservatives, formulated with *Mannheimia haemolytica* leukotoxin and lipopolysaccharide (LPS) supernatant antigens, as well as *Pasteurella multocida* type A bacterin. **Ovipast** aids in the prevention of sheep and goat pneumonia (Pasteurellosis).

DOSAGE FORM

Vaccine suspension for injection of a 1 ml dose via the subcutaneous route. A booster vaccination of 1 ml should be administered 3 to 4 weeks later. Annual revaccination is recommended thereafter.

STORAGE

Store in a refrigerator between 2 °C and 8 °C. Do not freeze. Protect from high temperatures and direct sunlight. Do not use after the expiry date printed on the packaging and container. Do not store any vials into which a needle or other device has been inserted, for future use.

WARNINGS

- Do not slaughter animals for human consumption within 21 days of vaccination.
- While no marked reaction should follow vaccination, a transient swelling may appear at the site of injection, and some animals may show a moderate rise of temperature for one or two days.
- Do not use in animal species other than those indicated.
- Do not administer to animals in poor or extremely poor body condition.
- KEEP THE PRODUCT OUT OF REACH OF CHILDREN, UNINFORMED PERSONS AND ANIMALS.
- Although this product has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice; notify the registration holder and the Registrar.

PRECAUTIONS

- Observe aseptic precautions. Ensure that all vaccination equipment (needles, syringes etc) is clean and sterile prior to use.
- Ensure that all equipment is kept clean and sterile during vaccination.
- Do not mix **Ovipast** with any other vaccine or medication.
- Use entire contents of vials when opened and do not store unused vaccine for future use.
- Do not use the contents of damaged vials.
- It is good vaccination practice when handling the vaccine to avoid contact with the eyes, mucous membranes, hands and clothing.
- Wash and disinfect hands with a disinfectant after vaccination.
- Used disposable vaccine equipment, unused vaccine etc. must be disposed of according to the National Environmental Waste Act, 2008 (Act 59 of 2008).

DIRECTIONS FOR USE

- Shake the bottle well before and during use with gentle end-over-end mixing.
- Inject sheep and goats subcutaneously on the side of the neck. DO NOT deviate from the recommended route and injection site location.
- Sheep and goats may be vaccinated from age of weaning and older.
- **Ovipast** can be administered to pregnant ewes to ensure passive antibody transfer to the lamb provided the lamb consumes sufficient quality colostrum within 1 day.
- It is essential to adhere to the vaccination programme to maintain a satisfactory immune response.
- Animals may therefore require a full vaccination program to be completed before the period of highest risk associated with respiratory pneumonia.

PRESENTATION

Ovipast, a light-yellow liquid with an off-white sediment if left to settle, is presented in a pack size of a 100 ml high density polyethylene (HDPE) vial of natural colour, capped with a gold-coloured aluminium cap, labelled and packed in a product-specific outer cardboard carton.

REGISTRATION HOLDER AND MANUFACTURER

Design Biologix CC
Reg. No. CK 1992/028856/23
442 Rigel Avenue South
Erasmusrand
Pretoria
0181
Tel: +27 (0) 12 349 2772
www.designbio.co.za

MARKETED AND DISTRIBUTED BY

Ceva Animal Health (Pty) Ltd
Reg. No. 1973/016009/07
Cnr 16th & Douglas Roads
Randjespark
Midrand
1685
Tel: +27 (0) 11 312 4088
Email: regulatory.za@ceva.com
www.ceva.co.za

DATE OF PUBLICATION OF THIS PACKAGE INSERT

16 August 2019



Ovipast

VIR SKAPE & BOKKE

Reg. Nr. G4356 (Wet 36 van 1947)

AANDUIDINGS

Ovipast is 'n multi-valente entstof met adjuvant, sonder antibiotika of preserveermiddels, geformuleer met *Mannheimia haemolytica* leukotoksien en lipopolisakkaried (LPS) bo-vloeistof antigene, asook die bakterien van *Pasteurella multocida* tipe A. **Ovipast** is 'n hulpmiddel in die voorkoming van skaap- en bok longontsteking (Pasteurellose).

ROETE VAN TOEDIENING

Entstof suspensie vir inspuiting met 'n 1 ml dosis via die onderhuidse roete. 'n Skraagdosie van 1 ml moet 3 tot 4 weke later toegedien word. Daar word daarna 'n jaarlikse her-inenting aanbeveel.

BERGINGSAAWYSINGS

Bêre in 'n yskas tussen 2 °C en 8 °C. Moenie vries nie. Beskerm teen hoë temperature en direkte sonlig. Moenie gebruik na die vervaldatum wat op die verpakking en houer gedruk is nie. Moenie houters waarin 'n naald of ander toestel aangebring is, vir toekomstige gebruik bêre nie.

WAARSKUWINGS

- Moenie diere binne 21 dae na inenting vir menslike verbruik slag nie.
- Normaalweg volg geen merkbare reaksie op inenting nie, alhoewel 'n kortstondige swelling op die plek van inenting kan voorkom, en sommige diere 'n matige styging in temperatuur vir een of twee dae kan toon.
- Moenie in spesies anders as die aangeduide spesies gebruik nie.
- Moet nie toedien aan diere in 'n swak of uiters swak liggaamskondisie nie.
- HOU DIE PRODUK BUITE BEREIK VAN KINDERS, ONINGELIGTE PERSONE EN DIERE.
- Alhoewel hierdie produk breedvoerig onder 'n wye verskeidenheid toestande getoets is, mag dit vaal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer en die Registrateur.

VOORSORGMATREËLS

- Let op aseptiese voorsorgmaatreëls. Maak seker dat alle inentingstoerusting (naalde, spuite, ens.) skoon en steriel is voor gebruik.
- Sorg dat alle toerusting tydens inenting skoon en steriel gehou word.
- Moet nie **Ovipast** met enige ander entstof of medikasie meng nie.
- Gebruik die hele inhoud van die houer wanneer dit oopgemaak word en moet nie ongebruikte entstof vir toekomstige gebruik bêre nie.
- Moenie die inhoud van 'n beskadigde houer gebruik nie.
- Dit is goeie inentingspraktyk om die entstof te hanteer om kontak met die oë, slymvliese, hande en klere te vermy.
- Was hande met 'n ontsmettingsmiddel na inenting.
- Gebruikte weggooibare toerusting, ongebruikte entstof, ens. moet weggedoen word volgens die

GEBRUIKSAANWYSINGS

- Meng die bottel goed voor en tydens gebruik.
- Spuit skape en bokke onderhuids aan die kant van die nek. MOENIE van die aanbevole roete en inspuitingsplek afwyk nie.
- Skape en bokke kan ingeënt word vanaf speenouderdom.
- Die entstof kan aan dragtige ooie toegedien word sodat passiewe teenliggame na die lam oorgedra word mits die lam binne 1 dag voldoende kwaliteit kolostrum inneem.
- Dit is noodsaaklik om die inentingsprogram te volg om 'n bevredigende immuunrespons te handhaaf.
- Dit word vereis dat die volledige inentingsprogram voltooi moet word voor die periode met die grootste risiko wat verband hou met respiratoriese longontsteking.

AANBIEDING

Ovipast, 'n liggeel vloeistof met wit sediment, word aangebied in 'n pakgrootte van 'n 100 ml hoëdigtheid-poliëteleen (HDPE) houer van natuurlike kleur, bedek met 'n goudkleurige aluminium doppie, geëtiketteer en verpak in 'n produksiespesifieke buitenste karton.

REGISTRASIEHOUER EN VERVAARDIGER

Design Biologix BK
Reg. Nr. CK 1992/028856/23
Rigellaan-Suid 442
Erasmusrand
Pretoria
0181
Tel: +27 (0) 12 349 2772
www.designbio.co.za

BEMARK EN VERSPREI DEUR

Ceva Animal Health (Edms) Bpk
Reg. Nr. 1973/016009/07
h/v 16de & Douglasweg
Randjespark
Midrand
1685
Tel: +27 (0) 11 312 4088
Epos: regulatory.za@ceva.com
www.ceva.co.za

DATUM VAN PUBLIKASIE VAN HIERDIE VOUBILJET

16 Augustus 2019

